

HOW TO USE...

ZOMACTON® 10mg
(somatropin) for Injection

Pocket Guide



ZOMACTON® (somatropin) for Injection Supplies



ZOMACTON 10 mg
(powder)



Alcohol swabs



Syringe and injection needle



Dry sterile gauze, cotton ball
or adhesive bandages



Pre-filled syringe with diluent
(liquid) and 25-gauge mixing needle



Sharps disposal container

ZOMACTON or diluent should not be used in patients with a known sensitivity to any of its ingredients.

STEP 1 Prepare

- Before you use ZOMACTON for the first time, talk to your healthcare provider about being properly trained on how to use it.
- ZOMACTON 10 mg is not ready for immediate use
- You must first mix (or reconstitute) the dry powder with the liquid included with your medication
- Mix 1 mL of pre-filled liquid with your medication
- Please note that the liquids are different for the 5 mg and 10 mg vials*—do NOT use the 5 mg liquid with 10 mg ZOMACTON or vice versa, unless otherwise directed by your doctor

Preparing ZOMACTON 10 mg for injection



- Place the supplies you will need on a clean, flat surface in a well-lit area
- Wash your hands thoroughly with soap and water
- Remove the syringe tip cap from the top of the pre-filled liquid syringe
- Attach the included 25-gauge mixing needle with needle cap

STEP 1

Prepare (Continued)

Diluting ZOMACTON® (somatropin) for Injection

- Only use the liquid that comes with the 10 mg ZOMACTON to mix



- Remove the hard plastic cap of the ZOMACTON vial
- Clean the top of the vial with an alcohol swab



- Remove the needle cap from the pre-filled liquid syringe and insert the needle into the center of the rubber stopper on the vial



- Slowly inject the liquid against the side of the vial to avoid creating air bubbles. **Do not** squirt the liquid directly onto the powder
- When all the liquid is in the vial, remove the needle
- Re-cap the needle and throw away the syringe in a sharps disposal container



Mixing ZOMACTON




- Hold the ZOMACTON vial between your hands and gently roll it (**never shake it**) until the mixture is clear. This may take a few seconds
- Do not** inject the medication if it is cloudy or has particles
- If air bubbles appear, let the mixture sit until they disappear
- Write the date you mixed ZOMACTON on the vial label. The 10 mg vial must be used within 28 days
- Store your **mixed** vial and all **unopened vials** of ZOMACTON in the refrigerator at 36°F to 46°F (2°C to 8°C). **Do not freeze**



STEP 2 Site Selection

ZOMACTON® (somatropin) for Injection can be injected into any of the following areas of the body (these sites should be rotated):



STEP 3 Inject



- Wash hands thoroughly with soap and water
- Check that the ZOMACTON vial is clear and the date of mixing is within 28 days
- Clean the top of the ZOMACTON vial with an alcohol swab. **Do not** touch the rubber stopper after cleaning



- Remove the needle cap from the injection syringe



- Slowly insert the needle into the center of the rubber stopper of the ZOMACTON vial



- Keeping the tip of the needle below the surface of the medication, turn the vial upside down
 - Gently pull back on the plunger until your prescribed amount of medication is in the syringe. (If you do not have enough medication for a full dose, reconstitute a new vial to make up the difference)

Contact your doctor if you notice any of the following:

- Lump, bruise, or redness at the injection site that doesn't go away
- Any sign of infection at the injection site (pus, redness, heat, or persistent pain)
- Severe, sharp pain or ache at injection site that does not go away
- Rash at injection site

STEP 3

Inject (Continued)



it means you've entered a blood vessel. **DO NOT** INJECT ZOMACTON; withdraw the needle, discard all supplies, and go back to step 1. Choose and clean a new injection site)

- If no blood appears, slowly push the plunger until the syringe is empty
- Quickly pull the needle straight out and apply pressure with a small dry sterile gauze pad or cotton ball, if needed. A drop of blood may appear. If it does, apply a small bandage. Throw away the needle and syringe in your sharps disposal container
- **Do not** share your syringes, needles, or vials with anyone else. You may give them or get an infection from them

zomacton.com

STEP 4

Store/Discard



- Once mixed the 10 mg vial must be used within 28 days
- Store the **mixed** ZOMACTON and all **unopened vials** in the refrigerator at 36°F to 46°F (2°C to 8°C). **Do not freeze**
- For safety reasons, it is important to properly store your medication and correctly dispose of used needles and syringes
- Remember, needles can only be used once, **do not** try to reuse the needle
- Dispose of needles, syringes, and vials in a sharps disposal container; **never dispose** of needles, syringes, and vials in your household trash or recycling bin
- Keep your sharps disposal container out of the reach of children
- When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container. There may be state or local laws about how you should throw away used needles and syringes. For more information about safe sharps disposal, and for specific information about sharps disposal in the state that you live in, go to the FDA's website at: <http://www.fda.gov/safesharpsdisposal>

ZOMACTON® 10mg
(somatropin) for Injection



Points of Interest

- You should not take ZOMACTON® (somatropin) for Injection if you are allergic to any of the ingredients in the medicine. ZOMACTON 10 mg vial is reconstituted with bacteriostatic water for injection containing metacresol. ZOMACTON or diluent should not be used in patients with a known sensitivity to any of its ingredients
- Try to inject ZOMACTON around the same time each day, in different areas of the body (see step 2)
- Keep track of injections and where you injected
- Don't worry if you miss a dose; just give the next scheduled one
- Use needles only once
- Never use expired medication (10 mg ZOMACTON expires 28 days after mixing even if stored properly at 36° to 46°F (2° to 8°C). **Do not** freeze)
- **Do not** use the 5 mg liquid with 10 mg ZOMACTON or vice versa unless otherwise directed by your doctor
- Always contact your healthcare provider if you have any questions or concerns about therapy with ZOMACTON




Designed to help patients and their families

- Assess insurance coverage
- Receive injection training
- Review copay or cash offers
- Ongoing patient, parent and/or caregiver support

Call the number below for more details.

1-844-944-ZOGO (9646)



ZOMACTON® 10mg
(somatropin) for Injection



ZOMACTON® 10mg
(somatropin) for Injection

ZOMACTON® is a registered trademark of Ferring B.V.
© 2023 Ferring B.V. All rights reserved. US-ZN-1900014

FERRING
PHARMACEUTICALS